

Singular Perturbations and Ion Channel Problems

Weishi Liu

Jilin University and University of Kansas

2016 International Conference on Singular Perturbation Theory
and its Applications

Hefei University, Hefei, Anhui, China

June 24-28, 2016

Abstract

Ion channel problems concern macroscopic properties of ionic flow through nano-scale ion channels. It is no coincidence that singularly perturbed systems serve as suitable models for analyzing these multi-scale problems. The general framework of singular perturbations often reveals special structures (idealized physical situations) of multi-scale phenomena and allows one to extract concrete information for specific problems. This is the case for the Poisson-Nernst-Plank (PNP) systems as primitive models for ionic flows.

In this talk, we will describe the geometric singular perturbation framework for an analysis of PNP systems and report a number of concrete results that are directly relevant to central topics of ion channel problems. The talk is based on works with several collaborators.

OUTLINE

Part I: Background, models, and a specific GSP

- Ion channel problems and Poisson-Nernst-Planck (PNP) models
- A general framework of GSP + Special structures of PNP
 - ⇒ Singular orbits involving ALL physical parameters of the problem

Part II: A number of specific applications

- Reversal (Nernst) potential and reversal permanent charge
- Effects of permanent charges and channel shape
- Ion size effects via PNP w/ Hard-Sphere potentials

Part I

1. Ion channel structures: shape and permanent charge

Ion channel functions: ionic flow and Poisson-Nernst-Planck models

2. A framework for analyzing PNP systems

- General theory of geometric singular perturbations (GSP)
- Special structures of PNP (most important ingredients for concrete information)
- Matching: yields (local) double-layers and brings (global) BC into picture
- Governing systems for singular orbits of BVP of PNP

1. Ion channel, ionic flow, PNP model




Figure 1: What Are Ion Channels: Shape and Permanent Charge

1.1. A quasi-one-dim PNP model for ionic flows of n types of ion species:

$$\text{Poisson: } \frac{1}{h(x)} \frac{d}{dx} \left(\epsilon^2 h(x) \frac{d\phi}{dx} \right) = -e \left(\sum z_s c_s + Q(x) \right),$$

$$\text{Nernst-Planck: } \frac{dJ_j}{dx} = 0, \quad -J_j = \frac{1}{k_B T} D_j h(x) c_j \frac{d\mu_j}{dx}.$$

$$\text{BV: } \phi(0) = \mathcal{V}, \quad c_j(0) = L_j; \quad \phi(1) = 0, \quad c_j(1) = R_j.$$

ϕ —electric potential, ϵ^2 —dielectric, $h(x)$ —area over x , $Q(x)$ —permanent charge

c_j — concentration, J_j — flux density, z_j — valence, D_j — diffusion constant,

Electrochemical potential: $\mu_j(\phi, \{c_i\}) = \mu_j^{id} + \mu_j^{ex}$:

Ideal component $\mu_j^{id} = z_j e \phi + k_B T \ln c_j$; Excess potential μ_j^{ex} for ion size.

A key quantity: **Current-Voltage (I-V) relation** $\mathcal{I} = \sum z_j J_j(\mathcal{V}; L, R)$.

1.2 A brief background

- Nernst; Planck (1890s):
 - * Nernst-Planck equation (study delayed due to lack of experimental data?)
 - * [Nernst equation for Reversal Potential \(Goldman-Hodgkin-Katz equation\)](#)
- Gouy-Chapmann (1910s); O. Stern (1924): [Double layer phenomena](#)
- Debye-Hückel; Lars Onsager (1920s):
 - * Electrolytic solutions based on [Poisson-Boltzmann approximations](#)
 - * Corrected to some extent by Lars Onsager when he was less than 22.
- Hodgkin-Huxley (1952a-e):
 - * “Voltage-Clamp” technique for recording action potentials in the squid giant axon ([single cell w/ a population of channels](#))
 - * Hodgkin-Huxley’s phenomenological model describes how action potentials in neurons are initiated and propagated
- Katz-Miledi, Neher-Sakmann (1970s): [Single-channel](#) recording of current
-:
 - * permeation, selectivity, gating, layering, charge inversion, conductivity, etc.

2. GSP for cPNP w/ piecewise constant $Q(x)$: [Liu JDE 09]

2.1. Reformulate BVP to a connecting problem (after a rescaling)

Introduce $u = \varepsilon \dot{\phi}$ and $w = x$. cPNP becomes, for $k = 1, 2, \dots, n$,

$$\begin{aligned}\varepsilon \dot{\phi} &= u, & \varepsilon \dot{u} &= - \sum_{s=1}^n z_s c_s - Q(w) - \varepsilon \frac{h'(w)}{h(w)} u, \\ \varepsilon \dot{c}_k &= -z_k c_k u - \varepsilon J_k h^{-1}(w), & \dot{J} &= 0, & \dot{w} &= 1.\end{aligned}$$

Associated to boundary conditions, introduce

$$\begin{aligned}B_L &= \{(\phi, u, C, J, w) \in \mathbb{R}^{2n+3} : \phi = \mathcal{V}, C = L, w = 0\}, \\ B_R &= \{(\phi, u, C, J, w) \in \mathbb{R}^{2n+3} : \phi = 0, C = R, w = 1\}.\end{aligned}$$

BVP \iff A connecting orbit from B_L to B_R .

2.2. Construction of Singular Orbits over $[0, 1]$.

Pre-assign the values of ϕ , c_k 's at jump point x_j of $Q(x)$ for $j = 1, 2, \dots, m - 1$,

$$\phi(x_j) = \phi^{[j]}, \quad c_k(x_j) = c_k^{[j]}, \quad k = 1, 2, \dots, n \quad (1)$$

with given $\phi^{[0]} = \mathcal{V}$ and $c_k^{[0]} = L_k$ at $x_0 = 0$, $\phi^{[m]} = 0$ and $c_k^{[m]} = R_k$ at $x_m = 1$, and introduce the set, for $j = 0, 1, \dots, m$,

$$B_j = \{(\phi, u, C, J, w) : \phi = \phi^{[j]}, C = C^{[j]}, w = x_j\}. \quad (2)$$

Two main steps for a construction of a singular orbits over $[0, 1]$

- Singular orbits on $[x_{j-1}, x_j]$ between B_{j-1} and B_j with $Q(x) = Q_j$.
- Matching them at jump points $x = x_j$'s to form a singular orbit on $[0, 1]$.



2.2.1. Singular orbit over $[x_{j-1}, x_j]$ between B_{j-1} and B_j with $Q(x) = Q_j$.

Each such an orbit will consist of two singular layers $\Gamma^{[j-1,r]}$ at $x = x_{j-1}$, and $\Gamma^{[j,l]}$ at $x = x_j$, and a regular layer Λ_j over the interval $[x_{j-1}, x_j]$.




Figure 3: Singular orbit over $[x_{j-1}, x_j]$

– Fast dynamics and bdry/internal layers.

The slow manifold is $\mathcal{Z}_j = \{u = 0, \sum_{s=1}^n z_s c_s + Q_j = 0\}$.

Note that $\dim \mathcal{Z}_j = 2n + 1 - \text{co-dim two}$.

In terms of the independent variable $\xi = x/\epsilon$, we obtain the fast system,

$$\begin{aligned}\phi' &= u, & u' &= -\sum_{s=1}^n z_s c_s - Q_j - \varepsilon \frac{h'(w)}{h(w)} u, \\ c'_k &= -z_k c_k u - \varepsilon J_k h^{-1}(w), & J' &= 0, & w' &= \varepsilon.\end{aligned}$$

The limiting fast system is, for $k = 1, 2, \dots, n$,

$$\begin{aligned}\phi' &= u, & u' &= -\sum_{s=1}^n z_s c_s - Q_j, \\ c'_k &= -z_k c_k u, & J' &= 0, & w' &= 0.\end{aligned}$$

Two e-values normal to \mathcal{Z}_j are $\pm \sqrt{\sum z_s^2 c_s} (\text{Debye length})^{-1} \implies \mathcal{Z}_j$ is NH.

Special structure of the limiting fast system:

Proposition. *The limiting fast system has a complete set of $(2n+2)$ first integrals given by, for $k = 1, 2, \dots, n$,*

$$G_k = \ln c_k + z_k \phi, \quad G_{n+1} = \frac{1}{2}u^2 - \sum_{s=1}^n c_s + Q_j \phi,$$

$$G_{n+1+k} = J_k \quad \text{and} \quad G_{2n+2} = w.$$

Consequences:

One can determine $u^{[j-1,+]}$ and $u^{[j,-]}$, and $\omega(B_{j-1})$ and $\alpha(B_j)$ up to J .

– Slow dynamics to connect $\omega(B_{j-1})$ and $\alpha(B_j)$.

Introduce $u = \varepsilon p$, $z_n c_n = -\sum_{s=1}^{n-1} z_s c_s - Q_j - \varepsilon q$.

In replacing u with p and c_n with q , **slow system becomes**, for $k = 1, \dots, n-1$,

$$\begin{aligned}\dot{\phi} &= p, & \varepsilon \dot{p} &= q - \varepsilon \frac{h'(w)}{h(w)} p, \\ \varepsilon \dot{q} &= \left(\sum_{s=1}^{n-1} (z_s - z_n) z_s c_s - z_n Q_j - \varepsilon z_n q \right) p + h^{-1}(w) \sum_{s=1}^n z_s J_s, \\ \dot{c}_k &= -z_k p c_k - J_k h^{-1}(w), & \dot{J} &= 0, & \dot{w} &= 1.\end{aligned}$$

When $\varepsilon = 0$, it is

$$\begin{aligned}\dot{\phi} &= p, & 0 &= q, \\ 0 &= \left(\sum_{s=1}^{n-1} (z_s - z_n) z_s c_s - z_n Q_j \right) p + h^{-1}(w) \sum_{s=1}^n z_s J_s, \\ \dot{c}_k &= -z_k p c_k - J_k h^{-1}(w), & \dot{J} &= 0, & \dot{w} &= 1.\end{aligned}$$

For this system, the slow manifold is

$$\mathcal{S}_j = \left\{ p = -\frac{h^{-1}(w) \sum_{s=1}^n z_s J_s}{\sum_{s=1}^{n-1} (z_s - z_n) z_s c_s - z_n Q_j}, q = 0 \right\}.$$

The limiting slow dynamics on \mathcal{S}_j is, with $\mathcal{I} = \sum_{s=1}^n z_s J_s$,

$$\begin{aligned}\dot{\phi} &= -\frac{h^{-1}(w)\mathcal{I}}{\sum_{s=1}^{n-1} (z_s - z_n) z_s c_s - z_n Q_j}, \\ \dot{c}_k &= \frac{h^{-1}(w)\mathcal{I}}{\sum_{s=1}^{n-1} (z_s - z_n) z_s c_s - z_n Q_j} z_k c_k - h^{-1}(w) J_k, \\ \dot{J} &= 0, \quad \dot{w} = 1.\end{aligned}$$

Special structure of limiting slow dynamics: On slow manifold \mathcal{S}_j ,

$$\sum_{s=1}^{n-1} z_s c_s + Q_j = -z_n c_n \implies \sum_{s=1}^{n-1} (z_s - z_n) z_s c_s - z_n Q_j = \sum_{s=1}^n z_s^2 c_s > 0.$$

Multiply $h(w) \sum_{s=1}^n z_s^2 c_s$ on the RHS to get

$$\begin{aligned} \frac{d}{d\tau}\phi &= -\mathcal{I}, & \frac{d}{d\tau}C &= D(J)C, & \sum_{s=1}^n z_s c_s + Q_j &= 0, \\ \frac{d}{d\tau}J &= 0, & \frac{d}{d\tau}w &= h(w) \sum_{s=1}^n z_s^2 c_s, \end{aligned}$$

where $D(J) = \Gamma - Jb^T$ with $\Gamma = \mathcal{I} \operatorname{diag}(z_1, \dots, z_n)$ and $b^T = (z_1^2, \dots, z_n^2)$.

Solving this system from $\omega(B_{j-1})$ to $\alpha(B_j)$, one gets $J^{[j]}$ over $[x_{j-1}, x_j]$.

2.2.2. Global matching: $u^{[j,-]} = u^{[j,+]}$ and $J^{[1]} = J^{[2]} = \dots = J^{[m]}$.

$m - 1 + n(m - 1) = (n + 1)(m - 1) =$ the number of preassigned unknowns.

The result gives the governing system for singular orbits of the BVP.

Theorem [L-Xu JDE 15]. For $Q = 0$ and general n , there is a unique solution.

From Eisenberg-L. 07 SIMA for $n = 2$ and three-piece-one-nonzero Q :

$$z_1 c_1^a e^{z_1(\phi^a - \phi^{a,m})} + z_2 c_2^a e^{z_2(\phi^a - \phi^{a,m})} + Q = 0,$$

$$z_1 c_1^b e^{z_1(\phi^b - \phi^{b,m})} + z_2 c_2^b e^{z_2(\phi^b - \phi^{b,m})} + Q = 0,$$

$$\frac{z_2 - z_1}{z_2} c_1^{a,l} = c_1^a e^{z_1(\phi^a - \phi^{a,m})} + c_2^a e^{z_2(\phi^a - \phi^{a,m})} + Q(\phi^a - \phi^{a,m}),$$

$$\frac{z_2 - z_1}{z_2} c_1^{b,r} = c_1^b e^{z_1(\phi^b - \phi^{b,m})} + c_2^b e^{z_2(\phi^b - \phi^{b,m})} + Q(\phi^b - \phi^{b,m}),$$

$$J_1 = \frac{c_1^L - c_1^{a,l}}{H(a)} \left(1 + \frac{z_1(\phi^L - \phi^{a,l})}{\ln c_1^L - \ln c_1^{a,l}} \right) = \frac{c_1^{b,r} - c_1^R}{H(1) - H(b)} \left(1 + \frac{z_1(\phi^{b,r} - \phi^R)}{\ln c_1^{b,r} - \ln c_1^R} \right),$$

$$J_2 = \frac{c_2^L - c_2^{a,l}}{H(a)} \left(1 + \frac{z_2(\phi^L - \phi^{a,l})}{\ln c_2^L - \ln c_2^{a,l}} \right) = \frac{c_2^{b,r} - c_2^R}{H(1) - H(b)} \left(1 + \frac{z_2(\phi^{b,r} - \phi^R)}{\ln c_2^{b,r} - \ln c_2^R} \right),$$

$$\phi^{b,m} = \phi^{a,m} - (z_1 J_1 + z_2 J_2) y,$$

$$c_1^{b,m} = e^{z_1 z_2 (J_1 + J_2) y} c_1^{a,m} - \frac{Q J_1}{z_1 (J_1 + J_2)} \left(1 - e^{z_1 z_2 (J_1 + J_2) y} \right),$$

$$J_1 + J_2 = - \frac{(z_1 - z_2)(c_1^{a,m} - c_1^{b,m}) + z_2 Q(\phi^{a,m} - \phi^{b,m})}{z_2 (H(b) - H(a))}.$$

Part II

1. Reversal potential **and** reversal permanent charge
2. **Effects of** permanent charges and channel geometry
3. Ion size **effects via PNP** w/ Hard-Sphere potentials

1. Reversal charge and potential: Eisenberg-L.-Xu (Nonlinearity 15)

- From NP:
$$J_k \int_0^1 \frac{1}{h(x)D_k c_k(x)} dx = \mu_k(0) - \mu_k(1) = \frac{e}{k_B T} z_k \mathcal{V}_0 + \ln \frac{L_k}{R_k}.$$

The **sign** of J_k is determined by bdry electrochemical potentials.

Permanent charges cannot do anything about the sign of J_k BUT do affect the magnitude of J_k .

Can $Q(x)$ change the **sign** of $\mathcal{I} = \sum z_k J_k$?.

- Reversal permanent charge Q is defined to be the one that makes $\mathcal{I} = 0$.
- Reversal potential \mathcal{V}_0 is defined to be the one that makes $\mathcal{I} = 0$.
- Consider $Q(x) = Q^*$ for $x \in [x_1, x_2]$ and $Q(x) = 0$ for $x \notin [x_1, x_2]$.

Let
$$g(V, \mathcal{V}_0) := \sum_{s=1}^n \frac{z_s(L_s e^{z_s \mathcal{V}_0} - R_s)}{1 - x_2 + x_1 e^{z_s \mathcal{V}_0} + (x_2 - x_1) e^{z_s V}}.$$

Theorem. (i) Fix \mathcal{V}_0 . If V^* is a *real* root of $g(V, \mathcal{V}_0) = 0$, then

$$Q^* = f(V, \mathcal{V}_0) := - \sum_{s=1}^n z_s e^{z_s(\mathcal{V}_0 - V^*)} \frac{[1 - x_2 + (x_2 - x_1) e^{z_s V^*}] L_s + x_1 R_s}{1 - x_2 + x_1 e^{z_s \mathcal{V}_0} + (x_2 - x_1) e^{z_s V^*}},$$

is a *reversal permanent charge*.

(ii) Fix Q^* . There is a *unique* solution (V, \mathcal{V}_0) of the system

$$g(V, \mathcal{V}_0) = 0 \quad \text{and} \quad f(V, \mathcal{V}_0) = Q^*,$$

and the corresponding \mathcal{V}_0 is the *reversal (Nernst) potential*.

Theorem. (i) For $n = 2$, \exists a reversal charge Q^* if and only if

$$(L_1 e^{z_1 \mathcal{V}_0} - R_1)(L_2 e^{z_2 \mathcal{V}_0} - R_2) > 0.$$

If exists, the reversal charge Q^* is *unique*.

(ii) For $n = 3$ with $z_1 = 1$, $z_2 = 2$ and $z_3 = -1$, and for some bdry conditions, there are *at least TWO* reversal permanent charges.

Theorem. For $n = 2$ with $z_1 = 1 = -z_2$ ($L_j = L$ and $R_j = R$).

For some (\mathcal{V}_0, L, R) , the reversal permanent charge Q^* exists and

$$J_1(0) > J_2(0) > J_1(Q^*) = J_2(Q^*).$$

[Somewhat counterintuitive, if not, nobody seems to know before.]

2. Effects of small $Q(x)$ and channel geometry: Ji-L.-Zhang SIAP 15

Consider $Q(x) = Q_0$ over $[x_1, x_2]$ w/ $n = 2$;

Electroneutrality: $z_1 L_1 = -z_2 L_2 = L$ and $z_1 R_1 = -z_2 R_2 = R$.

$$J_k(Q_0, \varepsilon) = J_{k0} + J_{k1}Q_0 + O(\varepsilon, Q_0^2).$$

2.1. Effects of channel geometry on fluxes of zeroth order in Q_0

$$J_{10} = \frac{L - R}{z_1 H(1)(\ln L - \ln R)} \mu_1^\delta, \quad J_{20} = \frac{R - L}{z_2 H(1)(\ln L - \ln R)} \mu_2^\delta;$$

$$\mu_k^\delta := \mu_k(0) - \mu_k(1) = z_k \mathcal{V}_0 + \ln L - \ln R \quad \text{and} \quad H(1) = \int_0^1 A^{-1}(x) dx.$$

J_{10} doesn't depend on 2nd ion species and J_{20} doesn't depend on 1st ion species.

Effects of channel geometry for zeroth order in Q_0 are simple.

2.2. Effects of $Q(x)$ and channel geometry on 1st order fluxes

$$J_{11} = \frac{A(\mu_2^\delta - z_2 BV)}{(z_1 - z_2)H(1)(\ln L - \ln R)^2} \mu_1^\delta,$$

$$J_{21} = \frac{A(\mu_1^\delta - z_1 BV)}{(z_2 - z_1)H(1)(\ln L - \ln R)^2} \mu_2^\delta,$$

where, in terms of $\alpha = H(x_1)/H(1)$ and $\beta = H(x_2)/H(1)$,

$$A = A(L, R) = -\frac{(\beta - \alpha)(L - R)^2}{((1 - \alpha)L + \alpha R)((1 - \beta)L + \beta R)(\ln L - \ln R)},$$

$$B = B(L, R) = \frac{\ln((1 - \beta)L + \beta R) - \ln((1 - \alpha)L + \alpha R)}{A}.$$

J_{11} depends on 2nd ion species and J_{20} depends on 1st ion species.

More detailed channel geometry presents in 1st order J_{k1} .

2.2.1. Channel geometry for optimal permanent charge effects on fluxes

Theorem. $|J_{11}|$ and $|J_{21}|$ attain their maximums for $(\alpha, \beta) = (0, 1)$.

Recall that $\alpha = H(x_1)/H(1)$, $\beta = H(x_2)/H(1)$, $H(x) = \int_0^x h^{-1}(s)ds$.

Striking Consequences

A short and narrow neck “ $>$ ” A long and wide neck.

Short and Narrow : $x_2 - x_1 \ll 1$ and $h(x)$ is much smaller for $x \in (x_1, x_2)$.

Long and Wide: $x_2 - x_1 \approx 1$ and $h(x)$ is more uniform with more charges

Ion channels prefer short and narrow necks.

2.2.2. Charge effects on fluxes of positively charged and negatively charged ions

For $t > 0$, set

$$\gamma(t) = \frac{t \ln t - t + 1}{(t - 1) \ln t}.$$

Lemma. For $t > 0$, $0 < \gamma(t) < 1$, $\gamma'(t) > 0$, $\gamma(t) + \gamma(1/t) = 1$,

$$\lim_{t \rightarrow 0} \gamma(t) = 0, \quad \lim_{t \rightarrow 1} \gamma(t) = 1/2, \quad \lim_{t \rightarrow \infty} \gamma(t) = 1.$$

Theorem. Let V_q^1 and V_q^2 be as

$$V_q^1 = V_q^1(L, R) = -\frac{\ln L - \ln R}{z_2(1 - B)} \quad \text{and} \quad V_q^2 = V_q^2(L, R) = -\frac{\ln L - \ln R}{z_1(1 - B)}.$$

Then, for $t = L/R > 1$, one has

(i) if $\alpha < \gamma(t)$ and $\beta \in (\alpha, \beta_1)$, then $V_q^1 < 0 < V_q^2$; and,

(i1) for $V \in (V_q^1, V_q^2)$, small positive Q_0 reduces both $|J_1|$ and $|J_2|$;

(i2) for $V < V_q^1$, small positive Q_0 strengthens $|J_1|$ but reduces $|J_2|$;

(i3) for $V > V_q^2$, small positive Q_0 reduces $|J_1|$ but strengthens $|J_2|$;

(ii) if either $\alpha < \gamma(t)$ and $\beta \in (\beta_1, 1)$ or $\alpha \geq \gamma(t)$, then $V_q^1 > 0 > V_q^2$; and,

(ii1) for $V \in (V_q^2, V_q^1)$, small positive Q_0 strengthens both $|J_1|$ and $|J_2|$;

(ii2) for $V > V_q^1$, small positive Q_0 strengthens $|J_1|$ but reduces $|J_2|$;

(ii3) for $V < V_q^2$, small positive Q_0 reduces $|J_1|$ but strengthens $|J_2|$.

SMALL positive permanent charge can do anything but strengthening the flux of positively charged ions while reducing that of negatively charged ions.

In general, for $k = 1, 2$ with $z_1 > 0 > z_2$, set $\lambda_k(Q, \varepsilon) = \frac{J_k(Q, \varepsilon)}{J_k(0, \varepsilon)}$.

Since Q does not change the sign of $J_k(Q, \varepsilon)$, one has $\lambda_k(Q, \varepsilon) > 0$.

If $\lambda_k(Q, \varepsilon) > 1$, then $|J_k(Q, \varepsilon)| > |J_k(0, \varepsilon)|$; that is,

the flux J_k is strengthened by the permanent charge Q .

Theorem. *For positive permanent charge $Q(x)$, one has, for $\varepsilon > 0$ small,*

$$\lambda_2(Q, \varepsilon) > \lambda_1(Q, \varepsilon) \quad \text{—} \quad \text{A Universal Effect}$$

and each of the following is possible:

$$(i) \lambda_2(Q, \varepsilon) > 1 > \lambda_1(Q, \varepsilon);$$

$$(ii) \lambda_2(Q, \varepsilon) > \lambda_1(Q, \varepsilon) > 1;$$

$$(iii) 1 > \lambda_2(Q, \varepsilon) > \lambda_1(Q, \varepsilon).$$

Corollary. Given a *positive permanent charge* $Q(x)$, it is possible that

- (i) the flux of *anion is enhanced* while that of *cation is reduced*;
- (ii) the fluxes of both anion and cation are enhanced and, in this case,

$$\frac{J_2(Q) - J_2(0)}{J_2(0)} > \frac{J_1(Q) - J_1(0)}{J_1(0)} > 0,$$

i.e, *the relative amount of flux increased* for anion is *greater than* that for cation;

- (iii) the fluxes of both anion and cation are reduced and, in this case,

$$0 < \frac{J_2(0) - J_2(Q)}{J_2(0)} < \frac{J_1(0) - J_1(Q)}{J_1(0)},$$

i.e., *the relative amount of flux reduced* for anion is *smaller than* that for cation.

3. PNP with hard-sphere potentials (HS)

3.1. Why do we care about ion sizes

Serious weakness of cPNP: treating $\text{Na}^+ = \text{K}^+$

In real world, $\text{Na}^+ \neq \text{K}^+$ significantly

Key difference: $\text{Na}^+ < \text{K}^+$ in ion size

Na^+ -channels v.s. K^+ -channels - (Protein Structure: MacKinnon 2003 Nobel)

Excess potential μ_i^{ex} accounts for finite size of ions
to distinguish ions with same valence but different sizes.

3.2. A one-dim **non-local** HS potential μ_j^{HS}

Percus-Yevick (70s) and Rosenfeld (93) model: (exact)

$$\mu_j^{HS} = \frac{\delta\Omega(\{c_i\})}{\delta c_j},$$

where

$$\Omega(\{c_i\}) = - \int n_0(x; \{c_i\}) \ln(1 - n_1(x; \{c_i\})) dx,$$
$$n_l(x; \{c_i\}) = \sum_i \int c_i(x') \omega_l^i(x - x') dx', \quad l = 0, 1$$
$$\omega_0^i(x) = \frac{\delta(x - r_i) + \delta(x + r_i)}{2}, \quad \omega_1^i(x) = \Theta(r_i - |x|).$$

δ : Dirac function; Θ : Heaviside function; r_i : radius of i th ions.

Statistical mechanics and geometric measurements of objects

3.3. A **local** HS potential μ_j^{LHS} for 3-dim

Bikerman's model (42):

$$\mu_j^{LHS}(x) = -\ln \left(1 - \frac{4\pi}{3} \sum_i r_i^3 c_i(x) \right) - \text{not ion specific.}$$

Many refined models

Boublik-Mansoori-Carnahan-Starling-Leland model (70-71):

Very accurate and more sophisticated, up to lowest order in radii,

$$\mu_j^{LHS}(x) = 8 \sum_i (r_j + r_i)^3 c_i(x) + O(r^6) - \text{ion specific.}$$

Quasi-one-dim'l PNP with HS potential

$$\frac{1}{h(x)} \frac{d}{dx} \left(\varepsilon^2 h(x) \frac{d\phi}{dx} \right) = - \sum_{s=1}^n z_s c_s - Q(x),$$
$$\frac{dJ_j}{dx} = 0, \quad -J_j = D_j h(x) \left(\frac{dc_j}{dx} + z_j c_j \frac{d\phi}{dx} + \underline{c_j \frac{d\mu_j^{HS}}{dx}} \right)$$

with the boundary conditions

$$\phi(0) = V, \quad c_j(0) = L_j; \quad \phi(1) = 0, \quad c_j(1) = R_j.$$

3.4. Electroneutrality, $n = 2$: $L := z_1 L_1 = -z_2 L_2$, $R := z_1 R_1 = -z_2 R_2$

Ji-L. JDDE 12 (for PNP w/ 1d nonlocal HS & $Q = 0$)

Lin-L.-Yi-Zhang SIADS 13 (for PNP w/ 1d local HS & $Q = 0$)

Let $r = r_1$ (ionic radius of 1st ion species) and $\lambda = r_2/r_1$, and let

$$V_c = \frac{(\lambda - 1)(L - R)(\ln L - \ln R)}{(z_1 \lambda - z_2)((L + R)(\ln L - \ln R) - 2(L - R))} \\ - \frac{(D_1 - D_2)(L + R)(\ln L - \ln R)^2}{(z_1 D_1 - z_2 D_2)((L + R)(\ln L - \ln R) - 2(L - R))};$$

$$V^c = \frac{(L - R)(\ln L - \ln R)}{z_1((L + R)(\ln L - \ln R) - 2(L - R))} \\ - \frac{(D_1 - D_2)(L + R)(\ln L - \ln R)^2}{(z_1 D_1 - z_2 D_2)((L + R)(\ln L - \ln R) - 2(L - R))}.$$

Theorem. [Size-Balance-Voltage V_c]

For $\varepsilon > 0$ small and $r > 0$ small,

(i) if $V > V_c$, then ion sizes enhance current: $I(V; \varepsilon, r) > I(V; \varepsilon, 0)$;

(ii) if $V < V_c$, then ion sizes reduce current: $I(V; \varepsilon, r) < I(V; \varepsilon, 0)$.

Theorem. [Size-Selectivity-Voltage V^c – independent of λ]

For $\varepsilon > 0$ small and $r > 0$ small,

(i) if $V > V^c$, the current I is increasing in λ

(smaller positive ion species is ‘preferred’, say, Na^+ over K^+);

(ii) if $V < V^c$, the current I is decreasing in λ

(larger positive ion species is ‘preferred’, say, K^+ over Na^+).

Scaling Properties in Boundary Concentrations

Write $I(V; \varepsilon, r) = I_0(V; \varepsilon) + I_1(V; \varepsilon)r + o(r)$.

$$I_0(V; 0) = I_0(V; L_j, R_j) - \text{point-charge contribution};$$

$$I_1(V; 0) = I_1(V; L_j, R_j) - \text{ion size component};$$

$$V_c = V_c(L_j, R_j), \quad V^c = V^c(L_j, R_j) - \text{two critical voltages}.$$

Theorem. [Scaling Laws in Bdry Concentrations]

(i) I_0 scales linearly in (L_j, R_j) : $I_0(V; \sigma L_j, \sigma R_j) = \sigma I_0(V; L_j, R_j)$;

(ii) I_1 scales quadratically in (L_j, R_j) : $I_1(V; \sigma L_j, \sigma R_j) = \sigma^2 I_1(V; L_j, R_j)$;

(iii) V_c and V^c scale invariantly in (L_j, R_j) : $V_c^c(\sigma L_j, \sigma R_j) = V_c^c(L_j, R_j)$.

Many other important applications of GSP to PNP to Ion channel problems !!!

Thank You !